

Shattered Shame States and their Repair

The John Bowlby Memorial Lecture

Saturday March 10, 2007

Judith Lewis Herman, M.D.

Final Draft: September 2007

Department of Psychiatry
Harvard Medical School

Victims of Violence Program
Department of Psychiatry
Cambridge Health Alliance
26 Central Street
Somerville, MA 02143

617 354-7547
herman.j@comcast.net

Introduction: Shame and the Attachment Bond

The primate relational systems for attachment, care-giving, mating, and social ranking, inclusion, exclusion and cooperation form a platform upon which complex human social life is built. Under ordinary conditions of peace, I would suggest that shame is one of the primary regulators of social relations. Fear is the primary regulator only in circumstances where social structures for maintaining peace have broken down and social relations are ruled by violence.

As the attachment system was initially conceptualized by Bowlby (1973), fear was considered the primary regulator. Bowlby described anxiety and anger as the infant's emotional responses to separation, and crying, following, and clinging as behavioral responses. As a good Darwinian, he saw the adaptive function of the attachment system in maintaining the infant's proximity to the caretaker, thus affording protection. This basic attachment system is common to human beings and other primates.

On the platform of this attachment bond are built the child's first internal working models of human intimacy. In primary attachment relationships the child learns to imagine other minds and to be in dialogue with beloved caretakers. Hennighausen and Lyons-Ruth (2005) propose that as humans have evolved "from biologic to dialogic" relational modes, the attachment system has been "partially displaced from its primate base." Emotional sharing and signaling become the primary mode for regulating security of attachment. The infant preferentially seeks out the caregiver who best knows her mind and is most attuned to her emotional signals. She also learns to imagine how others think of her, to become self-conscious.

While major disruptions in the attachment system produce fear, by second year of life the child reacts to more subtle disruptions with shame. Trevarthen (2005) speaks of “the feeling of shame in failure that threatens loss of relationship and hopeless isolation.” Schore (1998) conceptualizes shame as toddler’s response to a disappointed expectation of “sparkling-eyed pleasure” in the maternal gaze. Ordinarily child’s abashed signals elicit a caring response. The child learns that shame states do not signify complete disruption of attachment bond and that they can be regulated. Through repeated experiences of this kind the child and caretaker learn to negotiate emotional attunement and mutuality in their relationship.

Where no corrective relational process takes place, pathological variations in attachment system can develop. In particular, we see disorganized attachment where the primary attachment figure is a source of fear. I would argue that we also see disorganized attachment where the primary attachment figure is a source of unremitting shame. In this case the child is torn between need for emotional attunement and fear of rejection or ridicule. She forms an internal working model of relationship in which her basic needs are inherently shameful.

We are beginning to see the long-term effects of these “shattered states” in prospective longitudinal studies of high-risk children. Ogawa and associates (1997) in the Minnesota study found that, as Liotti (1999) has explained, disorganized attachment in infancy was strongly correlated with adolescent dissociation. They also found that having a “psychologically unavailable” caretaker in infancy (as rated by observers in home visits) predicted pathological dissociation in adolescence. A second longitudinal study by Lyons-Ruth (2003) at Harvard Medical School independently reached the same

conclusions. Both disorganized attachment behavior on the part of the child and “maternal disrupted communication” at 18 months separately predicted dissociation in adolescence.

To unpack what was meant by “maternal disrupted communication,” the raters distinguished three styles: hostile, withdrawn, and fearful. I would suggest that all three styles of maternal communication would be likely to produce chronic shame states: the hostile style through criticism and ridicule, the withdrawn and fearful styles through repeated rejection of the child’s bids for emotional connection.

Our knowledge of developmental trajectory of children with disorganized attachment is still rudimentary, but it appears that by age seven many of these children have essentially substituted care-giving or social ranking systems for the damaged attachment system, as a way of controlling proximity to a caregiver who does not care (Lyons-Ruth and Jacobvitz, 1999). This can also be conceptualized as an attempt to avoid the constant shame of unrequited love. Both the Minnesota and the Harvard longitudinal studies have also shown us that children who developed disorganized attachment in infancy have later difficulties with peer relationships. They have not learned to negotiate social cooperation or mutuality.

At the Victims of Violence Program at Cambridge Hospital, where I work, the majority of our adult patients report histories of abuse in childhood. Some were abused by their primary caretaker, but more commonly, the abuse was at the hands of someone else. Perhaps the abuser was someone in the family whom the primary caretaker valued more than she valued the child, or perhaps the abuser was an acquaintance outside of the family who had access to the child because the primary caretaker was not paying close

attention. In either case, the primary caretaker was not a source of fear, but she was “psychologically unavailable.” It is this absence, this breach in the primary attachment relationship which leaves our patients with the profound conviction that they are unlovable. In treatment, we find again and again that the core issue is shame. Our patients live in a state of chronic humiliation which profoundly distorts their view of self and others. I will speak later of how we try to address these shattered shame states in psychotherapy.

Liotti (2004) speaks of trauma, dissociation, and disorganized attachment as three strands of a single braid. I would like to add a fourth strand to the braid, by focusing on the role of shame in the development of traumatic disorders.

Characteristics of Shame:

Shame can be likened to fear in many respects. Like fear it is as “fast-track” physiologic response that in intense forms can overwhelm higher cortical functions. Like fear it is also a social signal, with characteristic facial and postural signs that can be recognized across cultures (Darwin, 1872; Izard, 1971). The gaze aversion, bowed head, and hiding behaviors of shame are similar to appeasement displays of social animals (Keltner & Harker, 1998), and may serve a similar social function among human beings. From an evolutionary point of view, shame may serve an adaptive function as a primary mechanism for regulating the individual’s relations both to primary attachment figures and to the social group (Izard, 1977; Gilbert & McGuire, 1998).

Like fear, shame is a biologically stressful experience. In a meta-analysis of 208 laboratory studies, Dickerson and Kemeny (2004) demonstrated that socially embarrassing test conditions (for example, public speaking) reliably produced elevated

cortisol and ACTH responses in human subjects. Perhaps because we have not found a reliable way to evoke shame in laboratory animals, however, understanding of the neurobiology of shame is rudimentary as compared to the extensive literature on fear. Schore (2003) proposes that shame is mediated by the parasympathetic nervous system and serves as a sudden “brake” on excited arousal states.

More than a century ago, Darwin (1872) described blushing as the most characteristic sign of shame, and questioned “how it has arisen that the consciousness that others are attending to our personal appearance should have led to the capillaries, especially those of the face, instantly becoming filled with blood.” This question remains unsolved. A more recent review article (Leary, Britt, Cutlip & Templeton, 1992) notes that, while some of the available evidence implicates the parasympathetic nervous system, “knowledge of the physiological basis of blushing is meager and clearly ripe for future research.”

The subjective experience of shame is of an initial shock and flooding with painful emotion. Shame is a relatively wordless state, in which speech and thought are inhibited. It is also an acutely self-conscious state; the person feels small, ridiculous and exposed. There is a wish to hide, characteristically expressed by covering the face with the hands. The person wishes to “sink through the floor” or “crawl in a hole and die.” Shame is always implicitly a relational experience. According to Lewis (1987b), one of the early pioneers in the study of shame, “Shame is one’s own vicarious experience of the other’s scorn. The self-in-the-eyes-of-the-other is the focus of awareness... The experience of shame often occurs in the form of imagery, of looking or being looked at. Shame may also be played out as an internal colloquy, in which the whole self is

condemned.” Thus shame represents a complex form of mental representation, in which the person is able to imagine the mind of another.

Developmental Origins of Shame

Developmentally, shame appears in the second year of life. Erikson (1950) formulates the central conflict of this developmental stage as “Autonomy vs. Shame and Doubt.” Properly speaking no toddler is autonomous; rather, one might formulate the toddler’s developmental task as learning to regulate body, affect, desire, and will in attunement with others. Positive resolution of the conflicts of this stage of life creates the foundation for healthy pride and mutuality in relationships, both self-respect and respect for others. Schore (2003) traces the origins of shame to the primary attachment relationship. Separations, which evoke fear and protest in normal toddlers, do not evoke shame; rather, shame can be seen in reunion interactions, when the toddler’s excitement is met with indifference or disapproval. To a certain extent, such experiences are inevitable and normal, since no caregiver can be empathically attuned to her child at all times, and sometimes the caretaker must chastise the child. However, under normal circumstances, the child’s shame reaction, like the appeasement displays of other primates, evokes a sympathetic response which in turn dispels the feeling of shame. The breach in attachment is thus repaired. Through repetition of this sequence, Schore postulates that the securely attached toddler learns the limits of the caregiver’s tolerance and also learns to self-soothe and regulate shame states.

Though shame and guilt are often spoken of interchangeably, and though both can be considered social or moral emotions, the two states are quite distinct. Whereas shame is focused on the global self in relation to others, guilt is focused on a specific action that

the person has committed. Shame is an acutely self-conscious state in which the self is “split,” imagining the self in the eyes of the other; by contrast, in guilt the self is unified. In shame, the self is passive; in guilt the self is active. Shame is an acutely painful and disorganizing emotion; guilt may be experienced without intense affect. Shame engenders a desire to hide, escape, or to lash out at the person in whose eyes one feels ashamed. By contrast, guilt engenders a desire to undo the offense, to make amends. Finally, shame is discharged in restored eye contact and shared, good-humored laughter, while guilt is discharged in an act of reparation (Lewis, 1987a).

The Social Functions of Shame

Originating in the primary attachment relationship, shame generalizes to become an emotion that serves to regulate peer relationships, social hierarchy, and all the basic forms of social life. Scheff and Retzinger (2000), building on the work of Lynd (1958), Goffman (1967), and Lewis (1971), describe shame as the “master emotion of everyday life.” In their conceptualization, shame is the “signal of trouble in a relationship.”

Shame, for example, serves to regulate social distance. People experience shame both if others are too distant, as in the extreme case of shunning or ostracism, and if others come too close, as in the extreme case where personal boundaries are violated.

Shame also mediates attunement to indices of social value or status. In its milder forms, shame is the result of social slights or ridicule. Mild experiences of shame are a part of ordinary social life. The everyday family of shame emotions includes shyness, self-consciousness, embarrassment, and feeling foolish or ridiculous. Through ordinary experiences of shame, individuals learn the boundaries of socially acceptable behavior.

In more extreme forms, shame is the reaction to being treated in a degrading manner. The extreme family of shame emotions includes humiliation, self-loathing, and feelings of defilement, disgrace, or dishonor. In hierarchical societies, according to Miller (1997), disgust and contempt are “emotions of status demarcation” that consign to lower status those against whom they are directed. Relationships of dominance and subordination are inherently shaming. The social signals of subordinate status (bowed head, lowered eyes) are ritualized expressions of shame. In slavery, the most extreme forms of social subordination, the enslaved person exists in a permanently dishonored status that Patterson (1982) describes as “social death.”

Extreme social subordination is found in relationships of coercive control: in modern-day slavery which takes the form of forced labor or prostitution (Bales, 2005), in political tyrannies, and in the private familial tyrannies of domestic violence and child abuse. Relationships of coercive control are established and perpetuated by an array of methods that are recognizable across cultures (Amnesty International, 1973). Among these methods, violence and threat of violence instill fear, while other commonly used methods, such as control of bodily functions, social isolation, and degradation, primarily evoke shame.

Extreme or catastrophic experiences of shame are a signal of profound relational disruptions or violations. When methods of coercive control are used within primary attachment relationships, as occurs in the case of child abuse, the developing child learns nothing of ordinary social shame. Rather, the child is overwhelmed with extreme shame states. Fonagy, Target, Gergely, Allen, & Bateman (2003) describe the shame of the abused child as “an intense and destructive sense of self-disgust, verging on self-hatred.”

They explain that “the shame concerns being treated as a physical object in the very context where special personal recognition is expected.”

Schore (2003) describes catastrophic shame states as “self-disorganizing.” Indeed, it is a characteristic of shame that it can feed upon itself. The shamed person feels ashamed of feeling ashamed, enraged, and ashamed of being enraged. Lewis (1990) describes these self-amplifying, disorganizing shame states as “feeling traps.” She proposes that when shame states can not be resolved, they are expressed as symptoms.

Shame as a Predictor of Posttraumatic Symptoms

Although the literature on this subject is sparse, three recent studies document an association between shame and posttraumatic symptoms. Andrews, Brewin, Rose & Kirk (2000) interviewed 157 victims of violent crime within one month of the incident and asked directly about shame experiences. At six month follow-up, shame was the only independent predictor of PTSD symptoms. Talbot, Talbot & Tu (2004), examined the relationship between shame-proneness and dissociation in a population of 99 hospitalized women with and without histories of childhood abuse. Shame-proneness was measured with a modification of the Differential Emotions Scale (DES-IV; Izard, Libero, Putnam & Haynes, 1993). Greater shame-proneness was associated with higher levels of dissociation, especially among women who had experienced sexual trauma early in their development. Interestingly, some women who had been abused in childhood were not particularly shame-prone and had dissociative scores within the normal range. The sources of resiliency in these women are not well understood and warrant further study.

Finally, Dutra, Callahan, Forman, Mendelsohn & Herman (2008, in press), in a study of 137 trauma survivors seeking outpatient treatment, measured self-reported

shame schemas using a modified version of the Young Schema Questionnaire (YSQ-S; Young & Brown, 1999). Shame schemas were significantly correlated with measures of PTSD and dissociation. Shame schemas were also specifically correlated with self-reported suicidal risk variables, including recent suicide attempts, current suicidal ideation, and current suicidal plans. These data would support the inference that posttraumatic shame states can be life-threatening.

Addressing Shame in Psychotherapy

Understanding that shame is a normal reaction to disrupted social bonds allows patients to emerge from the “feeling trap” in which they feel ashamed of being ashamed. According to Lewis (1981), addressing shame directly in the psychotherapy relationship facilitates therapeutic work, by normalizing shame reactions and by giving patients a relational framework for containing and understanding them. She writes: “Adopting the viewpoint that shame is a normal state which accompanies the breaking of affectional bonds allows shame to take its place as a universal, normal human state of being. Analyzing shame reactions in an atmosphere in which their natural function is taken for granted makes analytic work considerably easier.... Perhaps the greatest therapeutic advantage of viewing shame and guilt as affectional bond controls is the emphasis placed on the patients’ efforts to restore their lost attachments.”

The therapist calls attention to the patient’s shame reactions as they happen, noticing the bowed head and averted gaze. The therapist then invites the patient to move out of the shamed position, to lift her head, to make eye contact, and to experience the restorative empathic connection of the treatment relationship. As shame is relieved, often patient and therapist will spontaneously begin to laugh together. Retzinger (1987)

explains that shared laughter restores a sense of social connection: “Shame is a major aspect of the human condition. It serves a fundamental purpose, enabling human beings to monitor their own behavior in relation to others... When shame is too great, one feels alienated, disconnected from others, and alone in the world. Laughter serves to reconnect these severed ties, breaking the spiral of shame-rage... Without both shame and laughter, complex social life would be impossible.”

Numerous verbal, paralinguistic and nonverbal cues should alert the therapist to shame states. The vocabulary of shame is extensive: words such as “ridiculous, foolish, silly, idiotic, stupid, dumb, humiliated, disrespected, helpless, weak, inept, dependent, small, inferior, unworthy, worthless, trivial, shy, vulnerable, uncomfortable, or embarrassed” may indicate feelings of shame. Paralinguistic cues include confusion of thought, hesitation, soft speech, mumbling, silences, stammering, long pauses, rapid speech, or tensely laughed words. Nonverbal cues include hiding behavior such as covering all or parts of the face, gaze aversion, with eyes downcast or averted, hanging head, hunching shoulders, squirming, fidgeting, blushing, biting or licking the lips, biting the tongue, or false smiling (Retzinger, 1995).

Courtois (1988), in her description of therapeutic work with incest survivors, observes that shame may be difficult to address directly because of the way it affects the transference. The patient may have difficulty trusting evidence of her therapist’s positive regard, because she expects the therapist to feel the same contempt for her that she has for herself. It may be necessary for the therapist to challenge this distorted perception gently but directly. Shame also affects the countertransference; as Lewis

(1987b) explains, shame is a contagious emotion, and the therapist may avoid addressing shame directly because of her own discomfort.

Cloitre, Cohen & Koenen (2006), in their manual for treatment of survivors of childhood abuse, devote a chapter to the creation of narratives of shame. They write: “In the same way that narratives of fear must be titrated so that the client experiences mastery over fear rather than a reinstatement of it, so too narratives of shame should be titrated so that the client experiences dignity rather than humiliation in the telling.” These authors identify numerous reasons for telling about shameful events. They point out that shame perpetuates the bond with the perpetrator; as long as the patient guards her shameful secrets, she may feel that the perpetrator is the only person who knows her intimately. Disclosure in the context of the therapy relationship is a mastery experience that leads to greater self-knowledge, greater self-compassion and reduced feelings of alienation.

Patients with dissociative disorders have the additional burden of shame and secrecy about their illness itself. In their paper on treatment of dissociative disorders, Turkus and Kahler (2006) write that psychoeducation “helps to undo the stigmatization and shame associated with being ill. We have heard the words *insane*, *crazy* and *freak* many times from patients who are traumatized. In fact, patients on our trauma unit have requested that we change the group name to *psycheducation* to eliminate any implication of *psycho*.”

Because of the power imbalance between patient and therapist, and because the patient exposes her most intimate thoughts and feelings without reciprocity, the therapy relationship is to some degree inherently shaming. For this reason among others, group psychotherapy may be a particularly valuable treatment modality for traumatized people

(Herman & Schatzow, 1984; van der Kolk, 1987; Herman, 1992; Talbot et al, 1999; Mendelsohn, Zachary & Harney, 2007). The group members are peers who approach one another on a social plane of equality. Moreover, group members are in a position to give compassionate support as well as to receive it. Thus they can feel themselves to be of value to the group and deserving of the support they receive. The group becomes a little society within which members experience inclusion, cooperation and mutuality.

Group treatment must be structured so that group members titrate their exposure and learn to stay present rather than dissociating, both while describing their own experiences and while listening to others. This requires the group leaders to take an active stance, intervening when they notice a group member is disconnected, and modeling the kind of empathic feedback that group members can expect both to give and to receive. The resultant feeling of group acceptance and belonging is a powerful antidote to long-held feelings of shame and stigma (Herman & Schatzow, 1984).

Conclusion

If the thesis of this paper is correct, the role of shame in traumatic disorders and disorders of attachment should be a potentially fruitful area for further study. In particular, future research is needed to develop a fuller understanding of the neurophysiology of shame, to elucidate the role of shame in disorganized attachment and in posttraumatic symptom formation, and to explore the potentially therapeutic effects of addressing shame as a central issue in the treatment of trauma survivors.

References:

- Amnesty International (1973). Report on Torture. London: Duckworth.
- Andrews B., Brewin C.R., Rose, S. & Kirk, M. (2000). Predicting PTSD symptoms in victims of violent crime: The role of shame, anger, and childhood abuse. *Journal of Abnormal Psychology* 109: 69-73.
- Bales, K. (2005). *Understanding Global Slavery: A Reader*. Berkeley: University of California Press.
- Bowlby, J., (1973). *Attachment and Loss, Vol. 2: Separation: Anxiety and Anger*. New York: Basic Books.
- Cloitre M., Cohen, L.R. & Koenen, K.C. (2006). *Treating Survivors of Childhood Abuse: Psychotherapy for the Interrupted Life*. NY: Guilford. Quote p. 290.
- Courtois, C.A. (1988). *Healing the Incest Wound: Adult Survivors in Therapy*. NY: Norton.
- Darwin, C. (1872). *The Expression of the Emotions in Man and Animals*. London: John Murray. Quote p. 327.
- Dickerson, S.S. & Kemeny. M.E. (2004). Acute stressors and cortisol responses: A theoretical integration and synthesis of laboratory research. *Psychological Bulletin* 130:355-391.
- Dutra L., Callahan K., Forman E., Mendelsohn, M., & Herman, J.L. (2008, in press): Core schemas and suicidality in a chronically traumatized population. *Journal of Nervous and Mental Disease*.
- Erikson, E. H. (1950). *Childhood and Society*. New York: Norton.

- Fonagy, P., Target, M., Gergely, G., Allen, J.G., & Bateman, A.W. (2003). The developmental roots of Borderline Personality Disorder in early attachment relationships: A theory and some evidence. *Psychoanalytic Inquiry* 23: 412-459. Quote p. 445.
- Gilbert, P. & McGuire, M.T. (1998). Shame, status and social roles: Psychobiology and evolution. In P. Gilbert & B. Andrews (Eds.): *Shame: Interpersonal Behavior, Psychopathology and Culture*. NY Oxford University Press, 99-125.
- Goffman, E. (1967). *Interaction Ritual*. New York: Anchor
- Hennighausen K. H. & Lyons-Ruth, K., (2005). Disorganization of behavioral and attentional strategies toward primary attachment figures: From biologic to dialogic processes. In C.S, Carter, et.al (Eds.): *The 92nd Dahlem Workshop Report: Attachment and Bonding: A New Synthesis*. Cambridge, MA: MIT Press, 269-299.
- Herman J.L. (1992) *Trauma and Recovery*. New York: Basic.
- Herman J.L., & Schatzow, E. (1984). Time limited group therapy for women with a history of incest. *International Journal of Group Psychotherapy* 34: 605-615.
- Izard, C.E. (1971). *The Face of Emotion*. NY Appleton-Century-Crofts.
- Izard, C.E. (1977). *Human Emotions*. NY Plenum.
- Izard C.E., Libero, Z., Putnam, P., & Haynes, O.M. (1993). Stability of emotion experiences and their relations to traits of personality. *Journal of Personality and Social Psychology* 64: 847-860
- Keltner, D. & Harker, L.A. (1998). The forms and functions of the non-verbal signal of shame. In P Gilbert & B. Andrews (Eds.): *Shame: Interpersonal Behavior, Psychopathology and Culture*. NY Oxford University Press, 75-98.

- Leary, M.R., Britt, T.W., Cutlip, W.D. & Templeton J.L. (1992). Social blushing. *Psychological Bulletin* 112; 446-460.
- Lewis, H.B. (1971). *Shame and Guilt in Neurosis*. NY: International Universities Press.
- Lewis H.B. (1981). Shame and guilt in human nature. In S. Tuttmann, C. Kaye & M. Zimmerman (Eds.): *Object and Self: A Developmental Approach*. NY: International Universities Press, 235-265.
- Lewis H.B. (1987a). Shame and the narcissistic personality. In D. L. Nathanson (Ed.): *The Many Faces of Shame*. N.Y. Guilford, pp 93-132. Cf. Table on p. 113.
- Lewis H.B. (1987b). Introduction: Shame, the “sleeper” in psychopathology. In H.B. Lewis (Ed.) *The Role of Shame in Symptom Formation*. Hillsdale, NJ: Erlbaum. pp. 1-28. Quote on pp. 15, 18
- Lewis, H.B. (1990). Shame, repression, field dependence, and psychopathology. In J.L. Singer (Ed.). *Repression and Dissociation: Implications for Personality Theory, Psychopathology and Health*. University of Chicago Press, 233-257.
- Liotti, G., (1999). Disorganization of attachment as a model for understanding dissociative psychopathology. In J. Solomon & G. George (Eds.). *Disorganization of Attachment*. New York: Guilford, 291-317.
- Liotti, G (2004). Trauma, dissociation, and disorganized attachment: Three strands of a single braid. *Psychotherapy: Theory, Research, Practice, Training*. 41: 472-486.
- Lynd, H. (1958). *On Shame and the Search for Identity*. New York: Harcourt
- Lyons-Ruth, K. (2003). Dissociation and the parent-infant dialogue: A longitudinal perspective from attachment research. *Journal of the American Psychoanalytic Association* 51: 883-911.

Lyons-Ruth, K. & Jacobvitz, D., (1999). Attachment disorganization: Unresolved loss, relational violence, and lapses in behavioral and attentional strategies. In J. Cassidy & P. Shaver (Eds.): *Handbook of Attachment: Theory, Research, and Clinical Implications*. New York: Guilford, 520-554

Mendelsohn, M., Zachary, R.S., & Harney, P.A. (2007). Group therapy as an ecological bridge to new community for trauma survivors. In M.R. Harvey & P. Tummala-Narra (Eds.): *Sources and Expressions of Resiliency in Trauma Survivors*. New York: Haworth, 227-244.

Miller, W. (1997). *The Anatomy of Disgust*. Cambridge, MA: Harvard University Press.

Ogawa J. R., Sroufe, L. A., Weinfield, N. S., Carlson, E.A., & Egeland, B., (1997).

Development and the fragmented self: Longitudinal study of dissociative symptomatology in a nonclinical sample. *Development and Psychopathology* 9: 855-879.

Patterson, O. (1982). *Slavery and Social Death: A Comparative Study*. Cambridge, MA: Harvard University Press.

Retzinger, S.M. (1987): Resentment and laughter: Video studies of the shame-rage spiral. In H.B. Lewis (Ed.): *The Role of Shame in Symptom Formation*. Hillsdale NJ: Erlbaum, 151-181. Quote on p. 177.

Retzinger, S.M. (1995). Identifying shame and anger in discourse. *American Behavioral Scientist* 38:1104-1113.

Scheff, T.J. & Retzinger, S.M. (1997). Helen Block Lewis on shame: Appreciation and critique. In M.R. Lansky & A. Morrison (Eds). *The Widening Scope of Shame*. Hillsdale NJ: Analytic Press.

- Scheff, T.J. & Retzinger, S.M. (2000). Shame as the master emotion of everyday life. *Journal of Mundane Behavior* 1(3).
- Schore, A.N. (2003). *Affect Regulation and the Repair of the Self*. NY: Norton.
- Schore, A.N. (1998). Early shame experiences and infant brain development. In P. Gilbert & B. Andrews (Eds.): *Shame: Interpersonal Behavior, Psychopathology and Culture*. New York: Oxford, 57-77.
- Talbot, N., Houghtalen, R.P., Duberstein, P.R., Cox, C., Giles, D.E., & Wynne, L.C. (1999): Effects of group treatment for women with a history of childhood sexual abuse. *Psychiatric Services* 50: 686-692.
- Talbot, J., Talbot, N., Tu, X. (2004) Shame-proneness as a diathesis for dissociation in women with histories of childhood sexual abuse. *Journal of Traumatic Stress* 17:445-448.
- Trevarthen, C (2005). Stepping away from the mirror: Pride and shame in adventures of companionship: Reflections on the nature and emotional needs of infant intersubjectivity. In C.S. Carter, et. al (Eds.): *The 92nd Dahlem Workshop Report: Attachment and Bonding: A New Synthesis*. Cambridge, MA: MIT Press, 55-84.
- Turkus J.A, & Kahler J.A. (2006). Therapeutic interventions in the treatment of dissociative disorders. *Psychiatric Clinics of North America* 29: 245-262. Quote on page 246.
- Young, J.E. & Brown, G (1999). *Young schema questionnaire: Short form*. New York: Schema Therapy Institute.